




Socio-demographic patterning of objectively measured physical activity and sedentary behaviours in eight Latin American countries: Findings from the ELANS study

Gerson Luis de Moraes Ferrari, Irina Kovalskys, Mauro Fisberg, Georgina Gómez, Attilio Rigotti, Lilia Yadira Cortés Sanabria, Martha Cecilia Yépez García, Rossina Gabriella Pareja Torres, Marianella Herrera-Cuenca, Ioná Zalcman Zimberg, Viviana Guajardo, Michael Pratt, Carlos Cristi-Montero, Fernando Rodríguez-Rodríguez, Shaun Scholes, Carlos A. Celis-Morales, Jean-Philippe Chaput, Dirceu Solé & on behalf of the ELANS Study Group

To cite this article: Gerson Luis de Moraes Ferrari, Irina Kovalskys, Mauro Fisberg, Georgina Gómez, Attilio Rigotti, Lilia Yadira Cortés Sanabria, Martha Cecilia Yépez García, Rossina Gabriella Pareja Torres, Marianella Herrera-Cuenca, Ioná Zalcman Zimberg, Viviana Guajardo, Michael Pratt, Carlos Cristi-Montero, Fernando Rodríguez-Rodríguez, Shaun Scholes, Carlos A. Celis-Morales, Jean-Philippe Chaput, Dirceu Solé & on behalf of the ELANS Study Group (2019): Socio-demographic patterning of objectively measured physical activity and sedentary behaviours in eight Latin American countries: Findings from the ELANS study, *European Journal of Sport Science*, DOI: [10.1080/17461391.2019.1678671](https://doi.org/10.1080/17461391.2019.1678671)


To link to this article: <https://doi.org/10.1080/17461391.2019.1678671>

 [View supplementary material](#) 

 Accepted author version posted online: 11 Oct 2019.
Published online: 30 Oct 2019.

 [Submit your article to this journal](#) 


 Article views: 69

 [View related articles](#) 

 [View Crossmark data](#) 

ORIGINAL ARTICLE

Socio-demographic patterning of objectively measured physical activity and sedentary behaviours in eight Latin American countries: Findings from the ELANS study

GERSON LUIS DE MORAES FERRARI ^{1,2}, IRINA KOVALSKYS³, MAURO FISBERG^{2,4}, GEORGINA GÓMEZ⁵, ATTILIO RIGOTTI⁶, LILIA YADIRA CORTÉS SANABRIA⁷, MARTHA CECILIA YÉPEZ GARCÍA⁸, ROSSINA GABRIELLA PAREJA TORRES⁹, MARIANELLA HERRERA-CUENCA¹⁰, IONÁ ZALCMAN ZIMBERG¹¹, VIVIANA GUAJARDO³, MICHAEL PRATT¹², CARLOS CRISTI-MONTERO¹³, FERNANDO RODRÍGUEZ-RODRÍGUEZ¹³, SHAUN SCHOLES¹⁴, CARLOS A. CELIS-MORALES^{1,15}, JEAN-PHILIPPE CHAPUT¹⁶, DIRCEU SOLÉ², & ON BEHALF OF THE ELANS STUDY GROUP^{1†}

¹Centro de Investigación en Fisiología del Ejercicio - CIFE, Universidad Mayor, Santiago, Chile; ²Departamento de Pediatría da Universidade Federal de São Paulo, São Paulo, Brazil; ³Committee of Nutrition and Wellbeing, International Life Science Institute (ILSI-Argentina), Buenos Aires, Argentina; ⁴Instituto Pensi, Fundação José Luiz Egdio Setubal, Hospital Infantil Sabará, São Paulo, Brazil; ⁵Departamento de Bioquímica, Escuela de Medicina, Universidad de Costa Rica, San José, Costa Rica; ⁶Centro de Nutrición Molecular y Enfermedades Crónicas, Departamento de Nutrición, Diabetes y Metabolismo, Escuela de Medicina, Pontificia Universidad Católica, Santiago, Chile; ⁷Departamento de Nutrición y Bioquímica, Pontificia Universidad Javeriana, Bogotá, Colombia; ⁸Colégio de Ciencias de la Salud, Universidad San Francisco de Quito, Quito, Ecuador; ⁹Instituto de Investigación Nutricional, Lima, Peru; ¹⁰Centro de Estudios del Desarrollo, Universidad Central de Venezuela (CENDES-UCV)/Fundación Bengoa, Caracas, Venezuela; ¹¹Departamento de Psicobiología, Universidade Federal de São Paulo, São Paulo, Brazil; ¹²Institute for Public Health, University of California San Diego, La Jolla, CA, USA; ¹³IRyS Group, Physical Education School, Pontificia Universidad Católica de Valparaíso, Valparaíso, Chile; ¹⁴Institute of Epidemiology and Health Care, University College London, London, UK; ¹⁵British Heart Foundation Glasgow Cardiovascular Research Centre, Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, UK & ¹⁶Healthy Active Living and Obesity (HALO) Research Group, Children's Hospital of Eastern Ontario Research Institute, Ottawa, ON, Canada

ABSTRACT

Physical activity (PA) and sedentary behaviours (SB) are two independent risk factors for non-communicable diseases. However, there is a lack of objectively measured information on PA and SB in low- and middle-income countries. The aim of this study was to use objective data to characterise socio-demographic patterns of PA and SB in eight Latin American countries. 2732 participants (aged 15–65 years) from the Latin American Study of Nutrition and Health (ELANS) were included. PA and SB data were collected using accelerometers. Overall and country-specific average levels of time spent in PA and SB were compared by sex, age, socioeconomic and education level. Overall, the mean time spent in SB was 571.6 min/day, ranging from 553.8 min/day in Chile to 596.7 min/day in Peru. Average levels of light, moderate-to-vigorous physical activity (MVPA) and total PA were 311.1 min/day (95% CI: 307.7; 314.5), 34.9 min/day (95% CI: 34.0; 35.9) and 7531.2 MET-min/week (95% CI: 7450.4; 7611.9), respectively. MVPA and total PA were higher in men than women. The prevalence of physical inactivity was 40.6%, ranging from 26.9% (Chile) to 47% (Costa Rica and Venezuela). Women were more physically inactive than men (47.7% versus 33.0%). SB levels were highest among those with higher education; PA graded positively with socioeconomic level. Our findings can inform the planning of health policies and programmes designed to reduce levels of physical inactivity, as well as inform the local and cultural adaptation of these policies and programmes for implementation in Latin America.

[†]Membership of the ELANS Study Group is provided in the Acknowledgments section of the manuscript.

*Correspondence: Gerson Luis de Moraes Ferrari, Centro de Investigación en Fisiología del Ejercicio (CIFE), Universidad mayor - José Toribio Medina, 29 Estacion Central, Santiago, Chile. Email: gersonferrari08@yahoo.com.br

Highlights

- Worldwide studies of physical activity (PA) and sedentary behaviours (SB) have historically under-represented Latin American countries due to the lack of surveillance data.
- Across eight Latin American countries, the ELANS study collected data on PA and SB using an objective method (accelerometers) which we have analysed to quantify and characterise socio-demographic patterns.
- Over four-in-ten participants were physically inactive (40.6%); with a gender gap (47.7% women; 33.0% men); and striking differences between countries (47% Costa Rica and Venezuela; 26.9% Chile).
- In all countries, levels of moderate-to-vigorous physical activity (MVPA) were lowest, and levels of SB were highest, among participants in the higher education groups.
- Our findings on the unequal distribution of PA and SB increases the evidence base and can help to inform future intervention strategies in Latin America.

KEYWORDS: *Epidemiology, physical activity, sedentary behaviours, accelerometer, public health*

Introduction

Insufficient moderate-to-vigorous physical activity (MVPA) and excessive time spent in sedentary behaviours (SB) have emerged as key risk factors for cardiovascular disease worldwide, affecting not only high-income nations, but also low- and middle-income countries (LMICs) (Lee et al., 2012). More than three-in-four deaths due to chronic disease occur in LMICs (Lee et al., 2012). Therefore, in order to increase levels of physical activity (PA) and decrease the time spent in SB, researchers are joining efforts through a global call for action (Kohl et al., 2012; Sallis et al., 2016). However, regions such as Latin America continue to have scarce monitoring data on PA and SB.

Latin America countries has a unique structural, political, cultural and social environment (Salvo, Reis, Sarmiento, & Pratt, 2014). It is the most urbanised region worldwide (United Nations, 2012), with the rising prevalence of obesity and chronic diseases (Rivera, Pedraza, Martorell, & Gil, 2014). Latin America countries are characterised by high population density, pollution, and pronounced income inequality (United Nations, 2012). This makes Latin America a challenging but important region to gain knowledge about PA and SB to better inform intervention strategies.

Physical inactivity prevalence in Latin America is the highest reported worldwide (Guthold, Stevens, Riley, & Bull, 2018) and ranked fifth as a risk factor for mortality (GBD, 2017 Risk Factors Collaborators, 2018) according to the current global recommendations (World Health Organization, 2010) and US guidelines on PA for health (Physical Activity Guidelines Advisory Committee, 2018). The majority of population-based Latin American studies, conducted mainly in Brazil and Colombia, have used self-report methods to assess PA and SB, which have well-known limitations including problems of recall and reporting bias (Guthold, Stevens, Riley, & Bull, 2019; Silva et al., 2016).

Therefore, current population levels of PA and SB may be under- or over-estimated (Celis-Morales et al., 2012). Objective measures (e.g. accelerometry) can overcome some of these limitations and provide a more accurate estimate of the true levels of PA and SB (Hallal et al., 2012).

Use of accelerometers has increased in high-income countries (Compernelle et al., 2017; Kim, Im, & Choi, 2017). In contrast, relatively few studies have used accelerometers in LMICs, including Latin American countries, and few have implemented the same processing procedures to maximise comparability (Salvo et al., 2014). Only a few studies, the majority of which were completed in high-income countries, have used objectively measured PA data. Additionally, standard analysis protocols and cutoffs are uncommon for objective activity data (Guthold et al., 2019). Thus, the aim of this study was to use comparable accelerometer data to quantify and characterise socio-demographic patterns of PA and SB in Latin American countries.

Methods

Study design

The Latin American Study of Nutrition and Health (*Estudio Latinoamericano de Nutrición y Salud*; ELANS) is a cross-sectional, multi-national study conducted in eight countries (Argentina, Brazil, Chile, Colombia, Costa Rica, Ecuador, Peru and Venezuela) and aimed at investigating food and nutrient intake as well as nutritional and PA statuses of nationally representative samples from urban populations. The ELANS protocol was approved by the Institutional Review Board (#20140605) and registered at ClinicalTrials.gov (#NCT02226627). All participants provided informed consent/assent for participation in the ELANS study. Data were collected from September 2014 through February 2015. All the study sites followed the common protocol with all study personnel undergoing training and certification

in the data collection. More information on the ELANS is available elsewhere (Fisberg et al., 2016).

Participants

The recruitment of participants was conducted using a random complex, multistage sample design, stratified by conglomerates, with all regions of each country represented, and random selection of main cities within each region according to probability proportional to size method. The balanced sample was stratified by city, sex, age group and socioeconomic level for each country (Fisberg et al., 2016) (Supplementary Table A1). For the range of households within each secondary sampling unit, they were determined through systematic randomisation. The selection of the participants belonging to the domicile was made using 50% of the sample next birthday, and 50% last birthday. Thus, a total sample of 9218 (4809 [52.2%] women) participants (aged 15.0–65.0 years) was included in the ELANS study. In total, 92 cities were selected that participated in the study and the sampling size required for sufficient precision was calculated with a 95% confidence level and a maximum error of 3.5% and a survey design effect of 1.75 based on guidance from the National Center for Health Statistics (1996), and calculations of the minimum sample sizes required per sex, age group and socioeconomic level were performed for each country.

The sample with accelerometer data included 2372 participants aged 15–65 years, which represented 29.6% of the total ELANS cohort ($N=9218$) (Supplemental Table 1). Exclusion criteria included pregnant and lactating women, individuals with major physical or mental impairments, adolescents without assent and consent of a parent or legal guardian, individuals living in residential settings other than a household, and individuals unable to read (Fisberg et al., 2016). Considering puberty as a period in which occurs many biological and physiological changes, the inclusion of participants below 15 years of age would have required the determination of the pubertal stage, which is performed through the evaluation of the breasts, genitals and pubic hair (Keshteli et al., 2014). This was considered unfeasible in the current epidemiological study model. Thus, in order to ensure that puberty has been reached, only adolescents 15 years old and older were eligible to participate in the study.

Accelerometry assessment

The Actigraph GT3X accelerometer (Actigraph, Ft. Walton Beach, United States) was used to objectively monitor PA and SB. The accelerometer was worn on an elasticised belt at hip level on the right mid-axillary

line for seven days. Participants were asked to wear the device while awake and to remove it when sleeping, showering or swimming. Participants were encouraged to wear the accelerometer at least 12 h/day for at least seven days. The minimal amount of accelerometer data we considered acceptable for analytical purposes was five days (including at least one weekend day) with at least 10 h/day of wear time following the removal of sleep time (Colley, Connor Gorber, & Tremblay, 2010; Trost, Loprinzi, Moore, & Pfeiffer, 2011). After exclusion of the nocturnal sleep period time, waking nonwear time was defined as any sequence of at least 60 consecutive minutes of zero activity counts.

Data were processed using ActiLife software (V6.0; ActiGraph, Pensacola, FL). Data were collected at a sampling rate of 30 Hz and downloaded in epochs of 60 s (Brønd & Arvidsson, 2016).

SB was defined as time accumulated at ≤ 100 activity counts/min (1.5 METs), ≥ 101 –1951 activity counts/min for light PA (LPA; 1.6–2.9 METs), ≥ 1952 –5724 activity counts/min for moderate PA (MPA; 3.0–6.0 METs), ≥ 5725 activity counts/min for vigorous PA (VPA; >6.0 METs), and ≥ 1952 activity counts/min for MVPA (≥ 3.0 METs) (Dyrstad, Hansen, Holme, & Anderssen, 2014; Freedson, Melanson, & Sirard, 1998; Tremblay et al., 2017; Troiano et al., 2008). Reliability and validity of accelerometers have been documented extensively (Fisberg et al., 2016; Freedson, Lyden, Kozey-Keadle, & Staudenmayer, 2011). Participants were categorised as “meeting” or “not meeting” MVPA guidelines (World Health Organization, 2010).

Socio-demographic variables

Information about demographics including age, sex, years of education, race/ethnicity, and marital status was collected using standard questionnaires. Due to different classification systems across countries, we used three levels of classification based on the national statistics used in each country and included equivalent characteristics for all countries. Socioeconomic level data was divided into three strata (low, medium, high) based on the national indexes used in each country. Individuals self-reported their sex, age (15–29, 30–59, and ≥ 60 years), and education level (basic or lower [low], high school [middle], and university degree [high]). Detailed information can be found in a previous publication (Fisberg et al., 2016).

Statistical analysis

Descriptive statistics included mean and 95% confidence intervals (95% CI). Overall and country-

specific levels of PA and SB are presented by sex, age group, socioeconomic level, and educational level. Similar analyses were conducted for the prevalence of physical inactivity. Data analyses were performed with IBM SPSS, V.22. Analyses were weighted, with the weights calculated according to the socio-demographic characteristics, sex, socioeconomic level and region of each country (Fisberg et al., 2016). All tests of statistical significance were based on two-sided probability ($p < 0.05$).

Results

There were no significant differences between the participants wearing accelerometers and those who did not by sex ($p = 0.937$), socioeconomic level ($p = 0.501$) or educational level ($p = 0.235$). Participants wearing accelerometers were slightly older ($p = 0.018$). Accelerometers were worn on average for 15.3 h/day (Supplemental Table 2).

The proportion of women slightly exceeded that for men; 39.0% of participants were aged <30 years, 53.9% aged 30–59 years, and 7.1% aged ≥ 60 years. About half (51.2%) were classed as having a low socioeconomic level; 60.1% had a basic or lower educational level (Supplemental Table 3).

Overall, the mean SB and LPA were 571.6 min/day (95% CI: 567.1, 576.2) and 311.1 min/day (95% CI: 307.7, 314.5) respectively. Chile and Brazil had the lowest SB averages (Chile: 553.8 min/day; 95% CI: 540.1, 567.4; Brazil: 559.9 min/day; 95% CI: 550.0, 569.8) and the highest LPA (Chile: 324.4 min/day; 95% CI: 314.1, 334.7; Brazil: 321.2 min/day; 95% CI: 313.7, 328.7) (Supplemental Figure 1).

Overall, the mean MVPA was 34.9 min/day (95% CI: 34.0, 35.9); the average being highest in Chile (40.3 min/day; 95% CI: 37.4, 43.1) and lowest in Venezuela (31.2 min/day; 95% CI: 28.6, 33.7). In relation to total PA, the overall mean was 7531.2 MET-min/week (95% CI: 7450.4, 7611.9); the average being highest in Chile (7967.3 MET-min/week; 95% CI: 7723.8, 8210.8) and lowest in Costa Rica (7062.3 MET-min/week; 95% CI: 6808.3, 7316.2) (Supplemental Figure 1).

Overall, average levels of SB were higher in men than in women (582.7 min/day; 95% CI: 576.2, 589.2 versus 561.6 min/day; 95% CI: 555.3, 567.8). In relation to PA, women had higher values than men only in LPA (316.0 min/day; 95% CI: 311.3, 320.7 versus 305.7 min/day; 95% CI: 300.8, 310.7). MVPA was higher in men (42.1 min/day; 95% CI: 40.8, 43.4) than in women (28.3 min/day; 95% CI: 27.1, 29.6) overall and in each country. Average levels of total PA were slightly higher in

men than in women (7631.9 MET-min/week; 95% CI: 7515.1, 7748.1 versus 7439.2 MET-min/week; 95% CI: 7327.6, 7550.7) (Figure 1).

Average levels of SB varied by age. SB in persons aged <30 years was higher than in those aged 30–59 and ≥ 60 years in the total sample (591.5 min/day; 95% CI: 584.2, 598.9; 558.2 min/day; 95% CI: 552.2, 564.2; and 574.0 min/day; 95% CI: 557.2, 590.9 respectively) and in five of the eight countries (Argentina, Chile, Colombia, Costa Rica and Venezuela) (Figure 2).

Average levels of LPA (min/day) showed a positive gradient with age in all countries, with the exception of Colombia. Average levels of MVPA (min/day), with the exception of Colombia, showed a negative gradient with age (Figure 2).

A positive gradient between SB (min/day) and socioeconomic level was found in the full sample (Figure 3). The average levels of SB were 590.3 min/day (95% CI: 574.2, 606.5); 574.8 min/day (95% CI: 567.0, 582.7); and 562.7 min/day (95% CI: 556.3, 569.2) in the low, medium and high groups respectively. Average levels of LPA, MVPA and total PA were lower in the higher socioeconomic groups.

In the full sample, individuals with a higher education level spent more time in SB (594.2 min/day; 95% CI: 580.1, 608.3) than those in the middle (582.7 min/day; 95% CI: 574.4, 590.9) and low (562.3 min/day; 95% CI: 556.5, 568.1) education groups. Regarding the intensity of PA (MPA, VPA, MVPA and total), clear trends were not observed in all countries (Figure 4).

Overall, the prevalence of physical inactivity (<150 min/week in MVPA) was 40.6% (95% CI: 38.8, 42.5); ranging from 26.9% (Chile) to 47% (Costa Rica and Venezuela). Physical inactivity (>40%) was prevalent in Argentina (43.7%; 95% CI: 38.1, 49.5), Brazil (43.5%; 95% CI: 39.4, 47.7), Costa Rica (48.0%; 95% CI: 42.1, 53.9) and Venezuela (47.1%; 95% CI: 42.1, 52.3). For all countries, women (47.7%; 95% CI: 45.1, 50.3) were more likely to be inactive than men (33.0%; 95% CI: 30.4, 35.6) (Supplemental Figure 2).

Overall, physical inactivity prevalence was lower among those aged 30–59 years (36.7%; 95% CI: 34.2, 38.1) compared to those aged <30 years (45.3%; 95% CI: 42.2, 48.4) and those aged ≥ 60 years (47.4%; 95% CI: 40.4, 54.5). Results show a slightly higher percentage of physically inactive people among the participants with low socioeconomic level and education level. However, these trends were not observed in all countries. Those with a university degree (35.8% [95% CI: 30.4, 41.6]) had a lower prevalence of physical inactivity overall compared with those having a primary




Figure 1. Descriptive analysis (mean and 95% confidence interval) of sedentary behaviours, light, moderate, vigorous, moderate-to-vigorous physical activity (min/day) and total physical activity (MET-min/sem) of participants by sex from eight Latin America countries.




Figure 2. Descriptive analysis (mean and 95% confidence interval) of sedentary behaviours, light, moderate, vigorous, moderate-to-vigorous physical activity (min/day) and total physical activity (MET-min/sem) of participants by age group from eight Latin America countries.




Figure 3. Descriptive analysis (mean and 95% confidence interval) of sedentary behaviours, light, moderate, vigorous, moderate-to-vigorous physical activity (min/day) and total physical activity (MET-min/sem) of participants by socioeconomic level from eight Latin America countries.




Figure 4. Descriptive analysis (mean and 95% confidence interval) of sedentary behaviours, light, moderate, vigorous, moderate-to-vigorous physical activity (min/day) and total physical activity (MET-min/sem) of participants by education level from eight Latin America countries.

degree (44.7% [95% CI: 42.3; 47.1]) (Supplemental Figure 2).

Discussion

The aim of this study was to quantify and characterise socio-demographic patterns of PA and SB in eight Latin American countries. On average, participants spent 571.6 min/day in SB and 34.9 min/day in MVPA. MVPA was higher in men; and showed, with the exception of Colombia, a negative gradient with age. SB levels were highest among those with higher education. PA graded positively with socioeconomic level. Overall, just over four-in-ten participants were physically inactive.

This multicountry study quantified levels and characterised patterns of PA and SB in a diverse set of Latin American countries, using a comparable, reliable and validated instrument (Sasaki, John, & Freedson, 2011). Accelerometers were administered to 2732 participants, using a standard protocol. As such, this is the first study to report average levels of PA and SB – and the prevalence of physical inactivity – using nationally-representative samples from urban populations from Latin American countries based on objective measures. The only possible comparison was against previous reports describing PA at the regional level. Using accelerometers, some authors (Marsaux et al., 2016; Trost et al., 2011) have found different prevalence estimates of inactivity.

Guthold et al. (2018) compiled self-reported data from 358 surveys across 168 countries (1.9 million participants). Compared with the rest of the world, Latin American countries had high prevalence of insufficient PA (i.e. not achieving at least 150 min/week of MVPA). The highest levels were in Latin America and the Caribbean (39.1%), high-income Western countries (36.8%), high-income Asia Pacific (35.7%), south Asia (33.0%), central and eastern Europe (23.4%), east and southeast Asia (17.3%) and Oceania (16.3%).

Our own analyses based on accelerometer data show that the prevalence of insufficient PA varies greatly across the eight Latin American countries (Supplemental Figure 2); inactivity was about twice as high in Venezuela and Costa Rica compared to Chile. In high-income countries such as the United States and England, the transition towards more sedentary occupations and personal motorised transportation would explain, at least partially, the higher levels of inactivity (Ng & Popkin, 2012). In Latin American countries, levels of PA have been decreasing in both absolute and relative terms – reflecting large declines in occupational and domestic activity

and a small decline in active travel – while levels of SB have been increasing (Ng & Popkin, 2012). Therefore, it is necessary to implement municipal, regional and national policies to encourage active transportation, such as walking and cycling due to the rapid urban growth of some Latin American countries (United Nations, 2017), which has contributed to the high prevalence of physical inactivity in the region. Interventions should include better infrastructure for cycling and walking, improving safety and creating more opportunities for PA (World Health Organization, 2018). For example, México, Quito (Ecuador), Santiago (Chile), and Bogotá (Colombia) have Ciclovías, where roads are closed to motor vehicles on Sunday mornings and public holidays to permit people to contribute in PA (ÓDonovan, 2019) and studies of such initiatives in Bogotá have shown that users of Ciclovía programmes are more likely to comply with PA guidelines and have a higher quality of life (Montes et al., 2012).

Our study confirms findings (Althoff et al., 2017; Global Observatory for Physical Activity, 2018) of lower PA in women. Comparing sex differences in PA across countries is challenging, because patterns are strongly influenced by participation in different domains (which cannot be captured by accelerometers) and at different intensities of activity, which reflect the cultural and developmental contexts of each country. Mielke, da Silva, Kolbe-Alexander, and Brown (2018) shows that women spend less time in leisure activity than men. Offering more opportunities for safe and accessible leisure-time activity to women in order to increase their overall levels of activity would therefore help close the sex gap and achieve the 2025 global PA target of a 10% reduction in physical inactivity (Mielke et al., 2018).

Each country in our study had a mean level of SB higher than 9 h/day. Also in Latin America, Van Dyck et al. (2015) used accelerometers to evaluate SB in the International Physical Activity and the Environment Network study, showing means of 7.9 and 7.8 SB hours/day in Brazil and Colombia respectively. Objectively assessed SB from populations in Europe and the United States (National Health and Nutrition Examination Survey: NHANES) reported 7–8 h/day of SB (Compernelle et al., 2017; Kim et al., 2017). Most countries in the current study showed a socioeconomic and educational gradient in SB, with higher levels among the most educated. Further, SB was somewhat associated with the socioeconomic level. The reasons for higher average levels of SB in some countries cannot be determined from these descriptive data. Speculative reasons seem to point to adults with higher education and from high income countries having more sedentary jobs, being more

likely to use cars than active travel as a means of transport, and having more electronic entertainment and labor-saving devices at home (Bauman et al., 2009). Active transport is increasingly seen as an important opportunity to counteract the incidence of sedentary lifestyle diseases and is related to facilities and safety (Fishman, Böcker, & Helbich, 2015).

Some limitations of the present study are acknowledged. ELANS employed a cross-sectional design, precluding inferences about causality. The validity of cross-country comparisons may have been reduced to some extent by country-level variation in the questionnaire items on socioeconomic level (due to the legislative requirements or established local standard layouts). The cut-off point of counts/min for classifying the intensity of physical activity was the same for adolescents and adults and the accelerometers do not capture common activities such as cycling, resistance and static exercise, and carrying loads. Additionally, non-waterproof accelerometers cannot be used to assess water activities. Validation studies have compared accelerometry with questionnaires and found correlations of 0.33–0.40 between self-report and accelerometer (Dyrstad et al., 2014; Scholes et al., 2014). Questionnaires remain the most feasible method to assess PA levels at the population level owing in part to the expensive costs and high respondent burden associated with device-based methods in large-scale health examination surveys (Scholes, Bridges, Ng, & Mindell, 2016). Several questionnaires have been developed and standardised internationally for cross-study comparisons. Unfortunately, such standardised and validated questionnaires are often modified or translated without providing data on measurement properties and comparability of the altered version. By comparison, given their invariance to language and cultural differences, if standardised procedures existed, accelerometers would have greater potential for valid international comparisons. In view of the poor availability and representativeness of objectively measured PA data at present, Guthold et al. (2019) believe that national and global estimates of physical inactivity must be based on self-report data.

On the other hand, the strengths of this study include the large sample size, comparable data collection protocols, and the use of objective methods to assess PA and SB – use of these are rare in Latin America countries where most previous research has relied on self-reported instruments (de Sá et al., 2017). These types of objective assessments for PA and SB are rare for population health surveys and the best available evidence must be used to support and guide action to increase PA levels (Guthold et al., 2019). Our study is the first to evaluate PA

and SB patterns in Latin America using a standardised methodology across a consortium of several participating countries. This study thus provides a unique Latin American dataset that will enable wider cross-country comparisons and expand the existing literature.

Conclusion

The results from our cross-national study that objectively assessed PA and SB showed a high prevalence of physical inactivity and SB. Furthermore, we showed extensive variation of PA and SB levels across sociodemographic characteristics (sex, age, socioeconomic level and educational level). Our findings should help to inform the planning of health policies and programmes that are designed to reduce levels of physical inactivity, as well as the local and cultural adaptation of these policies and programmes for implementation in Latin America.

Acknowledgments

We would like to thank the following individuals at each of the participating sites who made substantial contributions to the ELANS: Luis A. Moreno, Beate Lloyd, Brenda Lynch, Mariela Jauregui, Alejandra Guidi, Luis Costa, and Regina Mara Fisberg. †The following are members of ELANS Study Group: Chairs: Mauro Fisberg, and Irina Kovalskys; Co-chair: Georgina Gómez Salas; Core Group members: Attilio Rigotti, Lilia Yadira Cortés Sanabria, Georgina Gómez Salas, Martha Cecilia Yépez García, Rossina Gabriella Pareja Torres, and Marianella Herrera-Cuenca; Steering Committee: Berthold Koletzko, Luis A. Moreno, Michael Pratt, and Katherine L. Tucker; Project Managers: Viviana Guajardo, and Ioná Zalcmán Zimberg; International Life Sciences Institute (ILSI)—Argentina: Irina Kovalskys, Viviana Guajardo, María Paz Amigo, Ximena Janezic, and Fernando Cardini; Universidad I Salud: Myriam Echeverry- Martin Langsman. Instituto Pensi-Hospital Infantil Sabara—Brazil: Mauro Fisberg, Ioná Zalcmán Zimberg, and Natasha Aparecida Grande de França; Pontificia Universidad Católica de Chile: Attilio Rigotti, Guadalupe Echeverría, Leslie Landaeta, and Óscar Castillo; Pontificia Universidad Javeriana-Colombia: Lilia Yadira Cortés Sanabria, Luz Nayibe Vargas, Luisa Fernanda Tobar, and Yuri Milena Castillo; Universidad de Costa Rica: Georgina Gómez Salas, Rafael Monge Rojas, and Anne Chinnock; Universidad San Francisco de Quito—Ecuador: Martha Cecilia Yépez García, Mónica Villar Cáceres, and María Belén Ocampo; Instituto de Investigación

Nutricional—Perú: Rossina Pareja Torres, María Reyna Liria, Krysty Meza, Mellisa Abad, and Mary Penny; Universidad Central de Venezuela: Mariannella Herrera-Cuenca, Maritza Landaeta, Betty Méndez, Maura Vasquez, Omaira Rivas, Carmen Meza, Servando Ruiz, Guillermo Ramirez, and Pablo Hernández; Statistical advisor: Alexandre D.P. Chiavegatto Filho; Accelerometry analysis: Priscila Bezerra Gonçalves, and Claudia Alberico; Physical activity advisor: Gerson Luis de Moraes Ferrari.

Funding

The ELANS was supported by a scientific grant from the Coca Cola Company, and support from the Ferrero, Instituto Pensi / Hospital Infantil Sabara, International Life Science Institute of Argentina, Universidad de Costa Rica, Pontificia Universidad Católica de Chile, Pontificia Universidad Javeriana, Universidad Central de Venezuela (CENDES-UCV)/Fundación Bengoa, Universidad San Francisco de Quito, and Instituto de Investigación Nutricional de Peru. The sponsors had no role in study design, in the collection, analyses, or interpretation of data, in the writing of the manuscript, and in the decision to publish the results.

Disclosure statement

No potential conflict of interest was reported by the authors.

Supplemental data

Supplemental data for this article can be accessed here doi:[10.1080/17461391.2019.1678671](https://doi.org/10.1080/17461391.2019.1678671).

ORCID

Gerson Luis de Moraes Ferrari  <http://orcid.org/0000-0003-3177-6576>

References

- Althoff, T., Sobic, R., Hicks, J. L., King, A. C., Delp, S. L., & Leskovec, J. (2017). Large-scale physical activity data reveal worldwide activity inequality. *Nature*, *547*(7663), 336–339.
- Bauman, A., Ainsworth, B. E., Bull, F., Craig, C. L., Hagstromer, M., Sallis, J. F., ... Sjostrom, M. (2009). Progress and pitfalls in the use of the international physical activity questionnaire (IPAQ) for adult physical activity surveillance. *Journal of Physical Activity and Health*, *6*(Suppl 1), S5–S8.
- Brønd, J. C., & Arvidsson, D. (2016). Sampling frequency affects the processing of Actigraph raw acceleration data to activity counts. *Journal of Applied Physiology*, *120*, 362–369.
- Celis-Morales, C. A., Perez-Bravo, F., Ibanez, L., Salas, C., Bailey, M. E., & Gill, J. M. (2012). Objective vs. self-reported physical activity and sedentary time: Effects of measurement method on relationships with risk biomarkers. *PLoS One*, *7*(5), e36345.
- Colley, R., Connor Gorber, S., & Tremblay, M. S. (2010). Quality control and data reduction procedures for accelerometry-derived measures of physical activity. *Health Reports*, *21*(1), 63–69.
- Compernelle, S., De Cocker, K., Mackenbach, J. D., Van Nassau, F., Lakerveld, J., Cardon, G., & De Bourdeaudhuij, I. (2017). Objectively measured physical environmental neighbourhood factors are not associated with accelerometer-determined total sedentary time in adults. *International Journal of Behavior Nutrition and Physical Activity*, *14*(1), 94.
- de Sá, T. H., Rezende, L. F. M., Borges, M. C., Nakamura, P. M., Anapolsky, S., Parra, D., et al. (2017). Prevalence of active transportation among adults in Latin America and the Caribbean: A systematic review of population based studies. *Revista Panamericana de Salud Publica*, *41*, e35.
- Dyrstad, S. M., Hansen, B. H., Holme, I. M., & Anderssen, S. A. (2014). Comparison of self-reported versus accelerometer-measured physical activity. *Medicine & Science in Sports and Exercise*, *46*(1), 99–106.
- Fisberg, M., Kovalskys, I., Gomez, G., Rigotti, A., Cortes, L. Y., Herrera-Cuenca, M., ... Guajardo, V. (2016). Latin American study of nutrition and health (ELANS): Rationale and study design. *BMC Public Health*, *16*(1), 93.
- Fishman, E., Böcker, L., & Helbich, M. (2015). Adult active transport in the Netherlands: An analysis of its contribution to physical activity requirements. *PLoS One*, *10*(4), e0121871.
- Freedson, P. S., Lyden, K., Kozey-Keadle, S., & Staudenmayer, J. (2011). Evaluation of artificial neural network algorithms for predicting METs and activity type from accelerometer data: Validation on an independent sample. *Journal of Applied Physiology*, *111*(6), 1804–1812.
- Freedson, P. S., Melanson, E., & Sirard, J. (1998). Calibration of the computer science and applications, Inc. Accelerometer. *Medicine & Science in Sports and Exercise*, *30*(5), 777–781.
- GBD 2017 Risk Factors Collaborators. (2018). Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: A systematic analysis for the global burden of disease study 2017. *The Lancet*, *392*, 1923–1994.
- Global Observatory for Physical Activity (GoPA). (2018). Country cards. Retrieved from <http://www.globalphysicalactivityobservatory.com/country-cards/>.
- Guthold, R., Stevens, G. A., Riley, L. M., & Bull, F. C. (2018). Worldwide trends in insufficient physical activity from 2001 to 2016: A pooled analysis of 358 population-based surveys with 1.9 million participants. *The Lancet Global Health*, *6*(10), e1077–e1086.
- Guthold, R., Stevens, G. A., Riley, L. M., & Bull, F. C. (2019). Accuracy and inequalities in physical activity research – authors’ reply. *The Lancet Global Health*, *7*(2), e187.
- Hallal, P. C., Andersen, L. B., Bull, F. C., Guthold, R., Haskell, W., Ekelund, U., & Lancet Physical Activity Series Working, G. (2012). Global physical activity levels: Surveillance progress, pitfalls, and prospects. *The Lancet*, *380*(9838), 247–257.
- Keshтели, A., Esmailzadeh, A., Rajaie, S., Askari, G., Feinle-Bisset, C., & Adibi, P. A. (2014). dish-based semi-quantitative food frequency questionnaire for assessment of dietary intakes in epidemiologic studies in Iran: Design and development. *International Journal of Preventive Medicine*, *5*(1), 29–36.
- Kim, J., Im, J. S., & Choi, Y. H. (2017). Objectively measured sedentary behavior and moderate-to-vigorous physical activity on the health-related quality of life in US adults: The national Health and Nutrition examination survey 2003–2006. *Quality of Life Research*, *26*(5), 1315–1326.
- Kohl, H. W., 3rd, Craig, C. L., Lambert, E. V., Inoue, S., Alkandari, J. R., Leetongin, G., Kahlmeier, S., Lancet Physical Activity Series Working, G., 2012. The pandemic of

- physical inactivity: Global action for public health. *The Lancet*, 380(9838), 294–305.
- Lee, I. M., Shiroma, E. J., Lobelo, F., Puska, P., Blair, S. N., Katzmarzyk, P. T., & Lancet Physical Activity Series Working, G., (2012). Effect of physical inactivity on major non-communicable diseases worldwide: An analysis of burden of disease and life expectancy. *The Lancet*, 380(9838), 219–229.
- Marsaux, C. F., Celis-Morales, C., Hoonhout, J., Claassen, A., Goris, A., Forster, H., ... Navas-Carretero, S. (2016). Objectively measured physical activity in European adults: Cross-sectional findings from the Food4Me study. *PLoS One*, 11(3), e0150902.
- Mielke, G. I., da Silva, I. C. M., Kolbe-Alexander, T. L., & Brown, W. J. (2018). Shifting the physical inactivity curve worldwide by closing the gender gap. *Sports Medicine*, 48(2), 481–489.
- Montes, F., Sarmiento, O. L., Zarama, R., Pratt, M., Wang, G., Jacoby, E., ... Ramos, M. (2012). Do health benefits outweigh the costs of mass recreational programs? An economic analysis of four Ciclovía programs. *Journal of Urban Health*, 89(1), 153–170.
- National Center for Health Statistics (NCHS). (1996). Analytic and reporting guidelines: The third national health and nutrition examination survey, NHANES III (1988–94). *Prevention*, 1–47.
- Ng, S. W., & Popkin, B. M. (2012). Time use and physical activity: A shift away from movement across the globe. *Obesity Reviews*, 13(8), 659–680.
- ÓDonovan, G. (2019). Accuracy and inequalities in physical activity research. *The Lancet Global Health*, 7(2), e186.
- Physical Activity Guidelines Advisory Committee. (2018). Physical activity guidelines advisory committee scientific report. Retrieved from <https://health.gov/paguidelines/second-edition/report/>.
- Rivera, J. A., Pedraza, L. S., Martorell, R., & Gil, A. (2014). Introduction to the double burden of undernutrition and excess weight in Latin America. *The American Journal of Clinical Nutrition*, 100(6), 1613S–1616S.
- Sallis, J. F., Bull, F., Guthold, R., Heath, G. W., Inoue, S., Kelly, P., ... Hallal, P. C. (2016). Progress in physical activity over the olympic quadrennium. *The Lancet*, 388(10051), 1325–1336.
- Salvo, D., Reis, R. S., Sarmiento, O. L., & Pratt, M. (2014). Overcoming the challenges of conducting physical activity and built environment research in Latin America: IPEN Latin America. *Preventive Medicine*, 69(Suppl 1), S86–S92.
- Sasaki, J. E., John, D., & Freedson, P. S. (2011). Validation and comparison of actiGraph activity monitors. *Journal of Science and Medicine in Sport*, 14(5), 411–416.
- Scholes, S., Bridges, S., Ng, F. L., & Mindell, J. S. (2016). Comparison of the physical activity and sedentary behaviour assessment questionnaire and the short-form international physical activity questionnaire: An analysis of health survey for England data. *PLoS One*, 11(3), e0151647.
- Scholes, S., Coombs, N., Pedisic, Z., Mindell, J. S., Bauman, A., & Rowlands, A. V. (2014). Age- and sex-specific criterion validity of the health survey for England physical activity and sedentary behavior assessment questionnaire as compared with accelerometry. *American Journal of Epidemiology*, 179(12), 1493–1502.
- Silva, L. J., Matsudo, V. K., Andrade, D. R., Azevedo, M., Ferrari, G. L., Oliveira, L. C., ... Matsudo, S. M. (2016). The prevalence of physical activity and its associated effects among students in the Sao Paulo public school network, Brazil. *Ciência & Saúde Coletiva*, 21(4), 1095–1103.
- Tremblay, M. S., Aubert, S., Barnes, J. D., Saunders, T. J., Carson, V., Cheung-Latimer, A. E., Chastin, S. F. M., Altenburg, T. M., Chinapaw, M. J. M., and on behalf of SBRN terminology consensus project participants. (2017). Sedentary behavior research network (SBRN) - terminology consensus project process and outcome. *International Journal of Behavioral Nutrition and Physical Activity*, 14(1), 75.
- Trojano, R. P., Berrigan, D., Dodd, K. W., Masse, L. C., Tilert, T., & McDowell, M. (2008). Physical activity in the United States measured by accelerometer. *Medicine & Science in Sports and Exercise*, 40(1), 181–188.
- Trost, S. G., Loprinzi, P. D., Moore, R., & Pfeiffer, K. A. (2011). Comparison of accelerometer cut points for predicting activity intensity in youth. *Medicine & Science in Sports and Exercise*, 43(7), 1360–1368.
- United Nations. (2012). *World urbanization prospects*. The 2011 Revision: Data tables and highlights 2011 rev ed.
- United Nations. (2017). *Department of economic and social affairs*. Population Division. World Urbanization Prospects, the 2014 revision. Retrieved from <https://esa.un.org/unpd/wup/>.
- Van Dyck, D., Cerin, E., De Bourdeaudhuij, I., Hinckson, E., Reis, R. S., Davey, R., ... Sallis, J. F. (2015). International study of objectively measured physical activity and sedentary time with body mass index and obesity: IPEN adult study. *International Journal Obesity (Lond)*, 39(2), 199–207.
- World Health Organization (WHO). (2010). *Global recommendations on physical activity for health*. Geneva: World Health Organization.
- World Health Organization (WHO). (2018). *Global action plan on physical activity 2018–2030. More active people for a healthier world*. Geneva: World Health Organization.