



IUCrData

ISSN: 2414-3146

[iucrdata.iucr.org/x](http://iucrdata.iucr.org/x)

## *N*-[3-(Prop-1-yn-1-yl)phenyl]benzenesulfonamide

Leslie W. Pineda and Jorge A. Cabezas

*IUCrData* (2019). **4**, x191176



IUCr Journals

CRYSTALLOGRAPHY JOURNALS ONLINE

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/4.0/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.





IUCrData

ISSN 2414-3146

# *N*-[3-(Prop-1-yn-1-yl)phenyl]benzenesulfonamide

Leslie W. Pineda<sup>a,b</sup> and Jorge A. Cabezas<sup>a\*</sup>

<sup>a</sup>Escuela de Química, Universidad de Costa Rica, 11501-2060, San José, Costa Rica, and <sup>b</sup>Centro de Electroquímica y Energía Química (CELEQ), Universidad de Costa Rica, 11501-2060, San José, Costa Rica. \*Correspondence e-mail: jorge.cabezas@ucr.ac.cr

Received 30 July 2019

Accepted 26 August 2019

Edited by W. T. A. Harrison, University of Aberdeen, Scotland


**Keywords:** crystal structure; antibacterial activity; N—H···O hydrogen bonding; propyne substituent; sulfonamide.

CCDC reference: 1887179


**Structural data:** full structural data are available from [iucrdata.iucr.org](http://iucrdata.iucr.org)

In the title sulfanilamide derivative, C<sub>15</sub>H<sub>13</sub>NO<sub>2</sub>S, which shows significant activity against *Staphylococcus aureus* and *Escherichia coli*, the dihedral angle between the planes of the aromatic rings is 62.15 (19)° and the four-coordinate S atom adopts an almost ideal tetrahedral geometry. In the crystal, N—H···O and C—H···O hydrogen bonds link the molecules into a three-dimensional network.

## 3D view



## Chemical scheme




## Structure description

In 1932, a drug called Prontosil was discovered by the pharmaceutical division of IG Farbenindustrie, an industrial conglomerate of German companies, including Bayer Company. It was found to be very successful treating several diseases in humans, provoked by *Staphylococcus* and *Streptococcus*. Prontosil was the first antibacterial drug, with life-saving capability, to be used systematically for the treatment of bacterial infections in the body. It belongs to a family of compounds called sulfa drugs or sulfonamides. In the 1940s and 1950s, most of the sulfa drugs were replaced by penicillin and other drugs, which proved to be more effective against more types of bacteria. However, nowadays, some sulfa drugs such as sulfamethoxazole, in combination with trimethoprim (co-trimoxazole), are still used extensively to inhibit the growth of bacteria that produce opportunistic infections in patients with AIDS, and bacterial infections such as pneumonia, bronchitis and infections of the urinary tract, ears and intestines (Brumfitt & Hamilton-Miller, 1993).

As part of our studies in this area we now report the synthesis of the title sulfanilamide derivative, **1**, and its crystal structure. This compound, has been found to be very effective against *Staphylococcus aureus* and *Escherichia coli*, and minimal inhibitory concentrations (MIC) of 12.5 µg ml<sup>-1</sup> and 25.0 µg ml<sup>-1</sup> have been obtained respectively (Cabezas & Arias, 2019).




OPEN ACCESS



**Figure 1**  
The title molecule with 50% probability ellipsoids.

The crystal structure of **1** has monoclinic symmetry with one molecule in the asymmetric unit: the molecular structure consists of a benzenesulfonamide fragment bound to a benzene ring bearing in its 3-position a propyne substituent (Fig. 1): the dihedral angle between the C1–C6 and C10–C15 benzene rings is 62.15 (19)°. The length of the carbon–carbon triple bond (C7≡C8) is 1.181 (5) Å, with the C7–C8–C9 and C8–C7–C1 angles being 178.8 (4) and 178.1 (4)°, respectively, which are slightly distorted from the expected linear geometry. The calculation of the angular structural index ( $\tau_4 = 0.94$ ; and  $\tau_4' = 0.90$ ) for the four-coordinate S1 atom, which binds to O1, O2, N1 and C10 from the benzene ring (Yang *et al.*, 2007; Okuniewski *et al.*, 2015; Rosiak *et al.*, 2018) indicates that it adopts an almost ideal tetrahedral geometry ( $\tau_4 = 0$  for an ideal square and 1 for an ideal tetrahedron). In the extended structure of **1**, weak N1–H1···O2, C4–H4···O1 and C6–H6···O1 hydrogen bonds are observed (Table 1, Fig. 2), leading to the formation of a three-dimensional network.



**Figure 2**  
Part of an [001] hydrogen-bonded chain with N–H···O2, C–H···O1 hydrogen bonds shown as green lines.

**Table 1**  
Hydrogen-bond geometry (Å, °).


| <i>D</i> –H··· <i>A</i>  | <i>D</i> –H | H··· <i>A</i> | <i>D</i> ··· <i>A</i> | <i>D</i> –H··· <i>A</i> |
|--------------------------|-------------|---------------|-----------------------|-------------------------|
| N1–H1···O2 <sup>i</sup>  | 0.88        | 2.55          | 2.984 (4)             | 111                     |
| C4–H4···O1 <sup>ii</sup> | 0.95        | 2.45          | 3.256 (4)             | 143                     |
| C6–H6···O1               | 0.95        | 2.39          | 2.955 (4)             | 118                     |

Symmetry codes: (i) *x*,  $-y + 1$ ,  $z + \frac{1}{2}$ ; (ii) *x*, *y*,  $z + 1$ .

**Table 2**  
Experimental details.

|   |   |
|---|---|
| Crystal data  |   |
| Chemical formula  | C <sub>15</sub> H <sub>13</sub> NO <sub>2</sub> S   |
| <i>M<sub>r</sub></i>  | 271.32  |
| Crystal system, space group   | Monoclinic, <i>Cc</i>   |
| Temperature (K)   | 100   |
| <i>a</i> , <i>b</i> , <i>c</i> (Å)  | 8.4596 (4), 24.9769 (13), 7.1310 (4)  |
| $\beta$ (°)   | 117.557 (2)   |
| <i>V</i> (Å <sup>3</sup> )  | 1335.80 (12)  |
| <i>Z</i>  | 4   |
| Radiation type  | Mo <i>K</i> $\alpha$  |
| $\mu$ (mm <sup>-1</sup> )   | 0.24  |
| Crystal size (mm)   | 0.35 × 0.20 × 0.15  |
| Data collection   |   |
| Diffractometer  | Bruker D8 Venture CCD   |
| Absorption correction   | Multi-scan ( <i>SADABS</i> ; Bruker, 2015)  |
| <i>T<sub>min</sub></i> , <i>T<sub>max</sub></i>   | 0.704, 0.746  |
| No. of measured, independent and observed [ <i>I</i> > 2 $\sigma$ ( <i>I</i> )] reflections                             | 9630, 3028, 2716  |
| <i>R<sub>int</sub></i>  | 0.037   |
| ( <i>sin</i> $\theta$ / $\lambda$ ) <sub>max</sub> (Å <sup>-1</sup> )   | 0.649   |
| Refinement  |   |
| <i>R</i> [ <i>F</i> <sup>2</sup> > 2 $\sigma$ ( <i>F</i> <sup>2</sup> )], <i>wR</i> ( <i>F</i> <sup>2</sup> ), <i>S</i> | 0.042, 0.089, 1.03  |
| No. of reflections  | 3028  |
| No. of parameters   | 173   |
| No. of restraints   | 2   |
| H-atom treatment  | H-atom parameters constrained   |
| $\Delta\rho_{max}$ , $\Delta\rho_{min}$ (e Å <sup>-3</sup> )  | 0.39, -0.47   |
| Absolute structure  | Flack <i>x</i> determined using 1163 quotients [( <i>I</i> <sup>+</sup> ) – ( <i>I</i> <sup>−</sup> )] / [( <i>I</i> <sup>+</sup> ) + ( <i>I</i> <sup>−</sup> )] (Parsons <i>et al.</i> , 2013) |
| Absolute structure parameter  | 0.02 (3)  |

Computer programs: *APEX3* and *SAINT* (Bruker, 2015), *SHELXT* (Sheldrick, 2015a), *SHELXL2014* (Sheldrick, 2015b), *Mercury* (Macrae *et al.*, 2006) and *publCIF* (Westrip, 2010).



**Figure 3**  
A synthetic scheme for the preparation of the title compound.

## Synthesis and crystallization

The title compound, **1**, was synthesized by treatment of 3-iodoaniline, **2**, with benzenesulfonyl chloride, **3**, in the presence of pyridine, at room temperature to obtain, after purification by column chromatography (ether:hexane, 40:60), iododisulfonamide, **4**, in 75% yield. This aromatic iodide **4**, was treated with propyne, under Sonogashira's reaction conditions (Sonogashira *et al.*, 1975), using CuI and (Ph<sub>3</sub>P)<sub>2</sub>PdCl<sub>2</sub> as catalysts, (Fig. 3). After purification by column chromatography, using a solvent mixture of hexane:ethyl acetate (75:25), compound **1** was isolated in 70% yield, and with an overall yield of 53%. The product was recrystallized from ethyl acetate solution at room temperature to result in light-yellow blocks of the title compound.

## Refinement

Crystal data, data collection and structure refinement are summarized in Table 2.

## Acknowledgements

CELEQ is thanked for supplying liquid nitrogen for the X-ray measurements.

## Funding information

Funding for this research was provided by: Vicerrectoría de Investigación, Universidad de Costa Rica (UCR); Escuela de Química (UCR).

## References

- Bruker (2015). *APEX3*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Brumfitt, W. & Hamilton-Miller, J. M. (1993). *J. Chemother.* **5**, 465–469.
- Cabezas, J. A. & Arias, M. L. (2019). *Int. J. Curr. Res.* **11**, 5224–5227.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Okuniewski, A., Rosiak, D., Chojnacki, J. & Becker, B. (2015). *Polyhedron*, **90**, 47–57.
- Rosiak, D., Okuniewski, A. & Chojnacki, J. (2018). *Polyhedron*, **146**, 35–41.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Sonogashira, K., Tohda, Y. & Hagihara, N. (1975). *Tetrahedron Lett.* **16**, 4467–4470.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Yang, L., Powell, D. R. & Houser, R. P. (2007). *Dalton Trans.* pp. 955–964.

## full crystallographic data

*IUCrData* (2019). 4, x191176 [https://doi.org/10.1107/S2414314619011763]

***N*-[3-(Prop-1-yn-1-yl)phenyl]benzenesulfonamide**

Leslie W. Pineda and Jorge A. Cabezas

***N*-[3-(Prop-1-yn-1-yl)phenyl]benzenesulfonamide***Crystal data*

$C_{15}H_{13}NO_2S$

$M_r = 271.32$

Monoclinic, *Cc*

$a = 8.4596$  (4) Å

$b = 24.9769$  (13) Å

$c = 7.1310$  (4) Å

$\beta = 117.557$  (2)°

$V = 1335.80$  (12) Å<sup>3</sup>

$Z = 4$

$F(000) = 568$

$D_x = 1.349$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 4995 reflections

$\theta = 2.8$ – $27.5$ °

$\mu = 0.24$  mm<sup>-1</sup>

$T = 100$  K

Block, clear light yellow

$0.35 \times 0.20 \times 0.15$  mm

*Data collection*

Bruker D8 Venture CCD

diffractometer

Radiation source: Incoatec Microsource

Mirrors monochromator

Detector resolution: 10.4167 pixels mm<sup>-1</sup>

$\omega$  scans

Absorption correction: multi-scan

(SADABS; Bruker, 2015)

$T_{\min} = 0.704$ ,  $T_{\max} = 0.746$

9630 measured reflections

3028 independent reflections

2716 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.037$

$\theta_{\max} = 27.5$ °,  $\theta_{\min} = 2.8$ °

$h = -10 \rightarrow 10$

$k = -32 \rightarrow 32$

$l = -9 \rightarrow 9$

*Refinement*

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.042$

$wR(F^2) = 0.089$

$S = 1.03$

3028 reflections

173 parameters

2 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0401P)^2 + 1.2876P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.39$  e Å<sup>-3</sup>

$\Delta\rho_{\min} = -0.47$  e Å<sup>-3</sup>

Absolute structure: Flack  $x$  determined using

1163 quotients  $[(I^-)-(I)]/[(I^+)+(I)]$  (Parsons et al., 2013)

Absolute structure parameter: 0.02 (3)

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** All hydrogen atoms were placed geometrically and refined using a riding-atom model approximation, with C—H = 0.95–1.00 Å, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . A rotating group model was used for the methyl groups.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

|     | <i>x</i>     | <i>y</i>     | <i>z</i>     | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|--------------|--------------|----------------------------------|
| S1  | 0.73168 (11) | 0.55417 (3)  | 0.33500 (12) | 0.01205 (19)                     |
| O1  | 0.7499 (3)   | 0.58926 (10) | 0.1883 (4)   | 0.0159 (5)                       |
| O2  | 0.7908 (3)   | 0.49971 (10) | 0.3513 (4)   | 0.0171 (6)                       |
| N1  | 0.8464 (4)   | 0.57903 (11) | 0.5727 (4)   | 0.0138 (6)                       |
| H1  | 0.9279       | 0.5593       | 0.6732       | 0.017*                           |
| C1  | 0.7946 (5)   | 0.72772 (15) | 0.5484 (6)   | 0.0188 (8)                       |
| C2  | 0.7594 (6)   | 0.73618 (15) | 0.7189 (6)   | 0.0243 (9)                       |
| H2  | 0.7403       | 0.7715       | 0.7534       | 0.029*                           |
| C3  | 0.7522 (6)   | 0.69340 (15) | 0.8383 (8)   | 0.0267 (9)                       |
| H3  | 0.7297       | 0.6994       | 0.9553       | 0.032*                           |
| C4  | 0.7781 (5)   | 0.64153 (15) | 0.7865 (6)   | 0.0176 (8)                       |
| H4  | 0.7705       | 0.612        | 0.8661       | 0.021*                           |
| C5  | 0.8149 (4)   | 0.63297 (13) | 0.6184 (5)   | 0.0134 (7)                       |
| C6  | 0.8256 (5)   | 0.67587 (14) | 0.5006 (5)   | 0.0156 (7)                       |
| H6  | 0.8538       | 0.6699       | 0.388        | 0.019*                           |
| C7  | 0.7989 (5)   | 0.77163 (15) | 0.4189 (6)   | 0.0216 (9)                       |
| C8  | 0.7988 (5)   | 0.80679 (16) | 0.3081 (6)   | 0.0232 (9)                       |
| C9  | 0.7969 (7)   | 0.85150 (17) | 0.1717 (8)   | 0.0345 (11)                      |
| H9A | 0.8463       | 0.8393       | 0.0788       | 0.052*                           |
| H9B | 0.8692       | 0.8811       | 0.26         | 0.052*                           |
| H9C | 0.6741       | 0.8637       | 0.0855       | 0.052*                           |
| C10 | 0.5068 (4)   | 0.55427 (14) | 0.2802 (5)   | 0.0138 (7)                       |
| C11 | 0.4331 (5)   | 0.50847 (14) | 0.3182 (5)   | 0.0159 (7)                       |
| H11 | 0.5029       | 0.477        | 0.3718       | 0.019*                           |
| C12 | 0.2569 (5)   | 0.50936 (17) | 0.2771 (6)   | 0.0207 (8)                       |
| H12 | 0.205        | 0.4782       | 0.3023       | 0.025*                           |
| C13 | 0.1549 (5)   | 0.55524 (16) | 0.1993 (6)   | 0.0202 (8)                       |
| H13 | 0.0343       | 0.5557       | 0.1737       | 0.024*                           |
| C14 | 0.2302 (5)   | 0.60067 (15) | 0.1587 (6)   | 0.0215 (8)                       |
| H14 | 0.1597       | 0.6319       | 0.1025       | 0.026*                           |
| C15 | 0.4069 (5)   | 0.60055 (14) | 0.1997 (6)   | 0.0168 (7)                       |
| H15 | 0.4588       | 0.6315       | 0.1733       | 0.02*                            |

*Atomic displacement parameters ( $\text{\AA}^2$ )*

|    | $U^{11}$   | $U^{22}$   | $U^{33}$   | $U^{12}$   | $U^{13}$   | $U^{23}$    |
|----|------------|------------|------------|------------|------------|-------------|
| S1 | 0.0130 (4) | 0.0114 (4) | 0.0127 (4) | 0.0013 (4) | 0.0067 (3) | −0.0001 (4) |

|     |             |             |             |              |             |              |
|-----|-------------|-------------|-------------|--------------|-------------|--------------|
| O1  | 0.0207 (14) | 0.0150 (12) | 0.0146 (12) | 0.0001 (10)  | 0.0103 (11) | 0.0000 (10)  |
| O2  | 0.0190 (13) | 0.0147 (12) | 0.0183 (13) | 0.0030 (10)  | 0.0094 (11) | -0.0010 (10) |
| N1  | 0.0139 (15) | 0.0131 (14) | 0.0117 (14) | 0.0025 (12)  | 0.0035 (12) | 0.0011 (11)  |
| C1  | 0.0196 (19) | 0.0142 (17) | 0.0189 (19) | -0.0017 (15) | 0.0057 (15) | 0.0006 (15)  |
| C2  | 0.037 (2)   | 0.0141 (19) | 0.022 (2)   | 0.0016 (17)  | 0.0139 (19) | -0.0035 (16) |
| C3  | 0.040 (3)   | 0.0230 (19) | 0.0225 (19) | 0.001 (2)    | 0.0188 (19) | -0.002 (2)   |
| C4  | 0.0222 (19) | 0.0162 (18) | 0.0174 (18) | -0.0003 (15) | 0.0119 (16) | 0.0039 (15)  |
| C5  | 0.0119 (17) | 0.0106 (17) | 0.0132 (17) | -0.0002 (13) | 0.0019 (14) | -0.0016 (13) |
| C6  | 0.0148 (17) | 0.0169 (18) | 0.0140 (17) | -0.0019 (14) | 0.0058 (14) | -0.0005 (14) |
| C7  | 0.026 (2)   | 0.0125 (19) | 0.025 (2)   | -0.0011 (16) | 0.0105 (17) | -0.0060 (16) |
| C8  | 0.028 (2)   | 0.017 (2)   | 0.025 (2)   | -0.0021 (17) | 0.0128 (19) | -0.0025 (18) |
| C9  | 0.043 (3)   | 0.024 (2)   | 0.039 (3)   | 0.001 (2)    | 0.021 (2)   | 0.011 (2)    |
| C10 | 0.0132 (15) | 0.0171 (17) | 0.0110 (16) | 0.0015 (15)  | 0.0054 (14) | -0.0030 (14) |
| C11 | 0.0180 (18) | 0.0158 (18) | 0.0129 (16) | -0.0009 (15) | 0.0062 (14) | 0.0010 (14)  |
| C12 | 0.0190 (19) | 0.028 (2)   | 0.0158 (18) | -0.0072 (16) | 0.0088 (15) | 0.0017 (16)  |
| C13 | 0.0142 (18) | 0.030 (2)   | 0.0172 (18) | -0.0017 (17) | 0.0078 (15) | -0.0075 (17) |
| C14 | 0.0182 (19) | 0.019 (2)   | 0.022 (2)   | 0.0015 (16)  | 0.0046 (16) | -0.0051 (16) |
| C15 | 0.0194 (19) | 0.0125 (17) | 0.0168 (17) | -0.0014 (14) | 0.0069 (15) | -0.0034 (14) |

*Geometric parameters (Å, °)*

|           |             |             |           |
|-----------|-------------|-------------|-----------|
| S1—O1     | 1.427 (3)   | C7—C8       | 1.181 (5) |
| S1—O2     | 1.435 (3)   | C8—C9       | 1.477 (5) |
| S1—N1     | 1.636 (3)   | C9—H9A      | 0.98      |
| S1—C10    | 1.756 (3)   | C9—H9B      | 0.98      |
| N1—C5     | 1.440 (4)   | C9—H9C      | 0.98      |
| N1—H1     | 0.88        | C10—C11     | 1.388 (5) |
| C1—C6     | 1.394 (5)   | C10—C15     | 1.390 (5) |
| C1—C2     | 1.396 (6)   | C11—C12     | 1.380 (5) |
| C1—C7     | 1.445 (5)   | C11—H11     | 0.95      |
| C2—C3     | 1.385 (6)   | C12—C13     | 1.387 (6) |
| C2—H2     | 0.95        | C12—H12     | 0.95      |
| C3—C4     | 1.392 (5)   | C13—C14     | 1.395 (5) |
| C3—H3     | 0.95        | C13—H13     | 0.95      |
| C4—C5     | 1.387 (5)   | C14—C15     | 1.384 (5) |
| C4—H4     | 0.95        | C14—H14     | 0.95      |
| C5—C6     | 1.390 (5)   | C15—H15     | 0.95      |
| C6—H6     | 0.95        |             |           |
| O1—S1—O2  | 119.34 (15) | C8—C7—C1    | 178.1 (4) |
| O1—S1—N1  | 108.24 (15) | C7—C8—C9    | 178.8 (4) |
| O2—S1—N1  | 104.96 (15) | C8—C9—H9A   | 109.5     |
| O1—S1—C10 | 108.01 (16) | C8—C9—H9B   | 109.5     |
| O2—S1—C10 | 108.67 (16) | H9A—C9—H9B  | 109.5     |
| N1—S1—C10 | 107.00 (15) | C8—C9—H9C   | 109.5     |
| C5—N1—S1  | 120.4 (2)   | H9A—C9—H9C  | 109.5     |
| C5—N1—H1  | 119.8       | H9B—C9—H9C  | 109.5     |
| S1—N1—H1  | 119.8       | C11—C10—C15 | 121.6 (3) |

|              |            |                 |            |
|--------------|------------|-----------------|------------|
| C6—C1—C2     | 119.5 (3)  | C11—C10—S1      | 119.6 (3)  |
| C6—C1—C7     | 119.1 (3)  | C15—C10—S1      | 118.8 (3)  |
| C2—C1—C7     | 121.4 (3)  | C12—C11—C10     | 118.9 (3)  |
| C3—C2—C1     | 120.4 (4)  | C12—C11—H11     | 120.5      |
| C3—C2—H2     | 119.8      | C10—C11—H11     | 120.5      |
| C1—C2—H2     | 119.8      | C11—C12—C13     | 120.6 (4)  |
| C2—C3—C4     | 119.9 (4)  | C11—C12—H12     | 119.7      |
| C2—C3—H3     | 120.1      | C13—C12—H12     | 119.7      |
| C4—C3—H3     | 120.1      | C12—C13—C14     | 119.7 (3)  |
| C5—C4—C3     | 119.9 (4)  | C12—C13—H13     | 120.1      |
| C5—C4—H4     | 120.1      | C14—C13—H13     | 120.1      |
| C3—C4—H4     | 120.1      | C15—C14—C13     | 120.4 (4)  |
| C4—C5—C6     | 120.4 (3)  | C15—C14—H14     | 119.8      |
| C4—C5—N1     | 118.6 (3)  | C13—C14—H14     | 119.8      |
| C6—C5—N1     | 120.9 (3)  | C14—C15—C10     | 118.7 (3)  |
| C5—C6—C1     | 119.8 (3)  | C14—C15—H15     | 120.7      |
| C5—C6—H6     | 120.1      | C10—C15—H15     | 120.7      |
| C1—C6—H6     | 120.1      |                 |            |
| O1—S1—N1—C5  | 54.7 (3)   | O1—S1—C10—C11   | 148.8 (3)  |
| O2—S1—N1—C5  | -176.8 (3) | O2—S1—C10—C11   | 18.0 (3)   |
| C10—S1—N1—C5 | -61.5 (3)  | N1—S1—C10—C11   | -94.9 (3)  |
| C6—C1—C2—C3  | -1.1 (6)   | O1—S1—C10—C15   | -31.3 (3)  |
| C7—C1—C2—C3  | 178.6 (4)  | O2—S1—C10—C15   | -162.1 (3) |
| C1—C2—C3—C4  | -0.8 (6)   | N1—S1—C10—C15   | 85.0 (3)   |
| C2—C3—C4—C5  | 1.4 (6)    | C15—C10—C11—C12 | -0.7 (5)   |
| C3—C4—C5—C6  | -0.2 (5)   | S1—C10—C11—C12  | 179.1 (3)  |
| C3—C4—C5—N1  | 178.1 (4)  | C10—C11—C12—C13 | -0.1 (5)   |
| S1—N1—C5—C4  | 126.3 (3)  | C11—C12—C13—C14 | 1.1 (6)    |
| S1—N1—C5—C6  | -55.4 (4)  | C12—C13—C14—C15 | -1.3 (6)   |
| C4—C5—C6—C1  | -1.6 (5)   | C13—C14—C15—C10 | 0.5 (5)    |
| N1—C5—C6—C1  | -179.9 (3) | C11—C10—C15—C14 | 0.5 (5)    |
| C2—C1—C6—C5  | 2.3 (5)    | S1—C10—C15—C14  | -179.3 (3) |
| C7—C1—C6—C5  | -177.5 (3) |                 |            |

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

| $D-H\cdots A$                   | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|---------------------------------|-------|-------------|-------------|---------------|
| N1—H1 $\cdots$ O2 <sup>i</sup>  | 0.88  | 2.55        | 2.984 (4)   | 111           |
| C4—H4 $\cdots$ O1 <sup>ii</sup> | 0.95  | 2.45        | 3.256 (4)   | 143           |
| C6—H6 $\cdots$ O1               | 0.95  | 2.39        | 2.955 (4)   | 118           |

Symmetry codes: (i)  $x, -y+1, z+1/2$ ; (ii)  $x, y, z+1$ .